

محاضرات التكامل

Integration



THOMAS'  
CALCULUS  
EARLY TRANSCENDENTALS

# Chapter 8

## Techniques of Integration

# 8.1

## Basic Integration Formulas

**TABLE 8.1** Basic integration formulas

1.  $\int du = u + C$
2.  $\int k du = ku + C$  (any number  $k$ )
3.  $\int (du + dv) = \int du + \int dv$
4.  $\int u^n du = \frac{u^{n+1}}{n+1} + C$  ( $n \neq -1$ )
5.  $\int \frac{du}{u} = \ln |u| + C$
6.  $\int \sin u du = -\cos u + C$
7.  $\int \cos u du = \sin u + C$
8.  $\int \sec^2 u du = \tan u + C$
9.  $\int \csc^2 u du = -\cot u + C$
10.  $\int \sec u \tan u du = \sec u + C$
11.  $\int \csc u \cot u du = -\csc u + C$
12.  $\int \tan u du = -\ln |\cos u| + C$   
 $= \ln |\sec u| + C$
13.  $\int \cot u du = \ln |\sin u| + C$   
 $= -\ln |\csc u| + C$
14.  $\int e^u du = e^u + C$
15.  $\int a^u du = \frac{a^u}{\ln a} + C$  ( $a > 0, a \neq 1$ )
16.  $\int \sinh u du = \cosh u + C$
17.  $\int \cosh u du = \sinh u + C$
18.  $\int \frac{du}{\sqrt{a^2 - u^2}} = \sin^{-1} \left( \frac{u}{a} \right) + C$
19.  $\int \frac{du}{a^2 + u^2} = \frac{1}{a} \tan^{-1} \left( \frac{u}{a} \right) + C$
20.  $\int \frac{du}{u\sqrt{u^2 - a^2}} = \frac{1}{a} \sec^{-1} \left| \frac{u}{a} \right| + C$
21.  $\int \frac{du}{\sqrt{a^2 + u^2}} = \sinh^{-1} \left( \frac{u}{a} \right) + C$  ( $a > 0$ )
22.  $\int \frac{du}{\sqrt{u^2 - a^2}} = \cosh^{-1} \left( \frac{u}{a} \right) + C$  ( $u > a > 0$ )

**TABLE 8.2** The secant and cosecant integrals

---

1.  $\int \sec u \, du = \ln |\sec u + \tan u| + C$

2.  $\int \csc u \, du = -\ln |\csc u + \cot u| + C$

## Procedures for Matching Integrals to Basic Formulas

### PROCEDURE

### EXAMPLE

Making a simplifying substitution

$$\frac{2x - 9}{\sqrt{x^2 - 9x + 1}} dx = \frac{du}{\sqrt{u}}$$

Completing the square

$$\sqrt{8x - x^2} = \sqrt{16 - (x - 4)^2}$$

Using a trigonometric identity

$$\begin{aligned}(\sec x + \tan x)^2 &= \sec^2 x + 2 \sec x \tan x + \tan^2 x \\ &= \sec^2 x + 2 \sec x \tan x \\ &\quad + (\sec^2 x - 1) \\ &= 2 \sec^2 x + 2 \sec x \tan x - 1\end{aligned}$$

Eliminating a square root

$$\sqrt{1 + \cos 4x} = \sqrt{2 \cos^2 2x} = \sqrt{2} |\cos 2x|$$

Reducing an improper fraction

$$\frac{3x^2 - 7x}{3x + 2} = x - 3 + \frac{6}{3x + 2}$$

Separating a fraction

$$\frac{3x + 2}{\sqrt{1 - x^2}} = \frac{3x}{\sqrt{1 - x^2}} + \frac{2}{\sqrt{1 - x^2}}$$

Multiplying by a form of 1

$$\begin{aligned}\sec x &= \sec x \cdot \frac{\sec x + \tan x}{\sec x + \tan x} \\ &= \frac{\sec^2 x + \sec x \tan x}{\sec x + \tan x}\end{aligned}$$

# 8.2

## Integration by Parts




$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx \quad (1)$$

### Integration by Parts Formula

$$\int u dv = uv - \int v du \quad (2)$$

### Integration by Parts Formula for Definite Integrals

$$\int_a^b f(x)g'(x) dx = f(x)g(x) \Big|_a^b - \int_a^b f'(x)g(x) dx \quad (3)$$



**FIGURE 8.1** The region in Example 6.

## EXAMPLE 8 Using Tabular Integration

Evaluate

$$\int x^3 \sin x \, dx.$$

**Solution** With  $f(x) = x^3$  and  $g(x) = \sin x$ , we list:

$f(x)$ and its derivatives		$g(x)$ and its integrals
$x^3$	(+)	$\sin x$
$3x^2$	(-)	$-\cos x$
$6x$	(+)	$-\sin x$
$6$	(-)	$\cos x$
$0$		$\sin x$

Again we combine the products of the functions connected by the arrows according to the operation signs above the arrows to obtain

$$\int x^3 \sin x \, dx = -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x + C.$$

# 8.3

## Integration of Rational Functions by Partial Fractions

## Method of Partial Fractions ( $f(x)/g(x)$ Proper)

1. Let  $x - r$  be a linear factor of  $g(x)$ . Suppose that  $(x - r)^m$  is the highest power of  $x - r$  that divides  $g(x)$ . Then, to this factor, assign the sum of the  $m$  partial fractions:

$$\frac{A_1}{x - r} + \frac{A_2}{(x - r)^2} + \cdots + \frac{A_m}{(x - r)^m}.$$

Do this for each distinct linear factor of  $g(x)$ .

2. Let  $x^2 + px + q$  be a quadratic factor of  $g(x)$ . Suppose that  $(x^2 + px + q)^n$  is the highest power of this factor that divides  $g(x)$ . Then, to this factor, assign the sum of the  $n$  partial fractions:

$$\frac{B_1x + C_1}{x^2 + px + q} + \frac{B_2x + C_2}{(x^2 + px + q)^2} + \cdots + \frac{B_nx + C_n}{(x^2 + px + q)^n}.$$

Do this for each distinct quadratic factor of  $g(x)$  that cannot be factored into linear factors with real coefficients.

3. Set the original fraction  $f(x)/g(x)$  equal to the sum of all these partial fractions. Clear the resulting equation of fractions and arrange the terms in decreasing powers of  $x$ .
4. Equate the coefficients of corresponding powers of  $x$  and solve the resulting equations for the undetermined coefficients.

## Heaviside Method

1. Write the quotient with  $g(x)$  factored:

$$\frac{f(x)}{g(x)} = \frac{f(x)}{(x - r_1)(x - r_2) \cdots (x - r_n)}.$$

2. Cover the factors  $(x - r_i)$  of  $g(x)$  one at a time, each time replacing all the uncovered  $x$ 's by the number  $r_i$ . This gives a number  $A_i$  for each root  $r_i$ :

$$A_1 = \frac{f(r_1)}{(r_1 - r_2) \cdots (r_1 - r_n)}$$

$$A_2 = \frac{f(r_2)}{(r_2 - r_1)(r_2 - r_3) \cdots (r_2 - r_n)}$$

⋮

$$A_n = \frac{f(r_n)}{(r_n - r_1)(r_n - r_2) \cdots (r_n - r_{n-1})}.$$

3. Write the partial-fraction expansion of  $f(x)/g(x)$  as

$$\frac{f(x)}{g(x)} = \frac{A_1}{(x - r_1)} + \frac{A_2}{(x - r_2)} + \cdots + \frac{A_n}{(x - r_n)}.$$

# 8.4

## Trigonometric Integrals

## Products of Powers of Sines and Cosines

We begin with integrals of the form:

$$\int \sin^m x \cos^n x dx,$$

where  $m$  and  $n$  are nonnegative integers (positive or zero). We can divide the work into three cases.

**Case 1** If  $m$  is odd, we write  $m$  as  $2k + 1$  and use the identity  $\sin^2 x = 1 - \cos^2 x$  to obtain

$$\sin^m x = \sin^{2k+1} x = (\sin^2 x)^k \sin x = (1 - \cos^2 x)^k \sin x. \quad (1)$$

Then we combine the single  $\sin x$  with  $dx$  in the integral and set  $\sin x dx$  equal to  $-d(\cos x)$ .

**Case 2** If  $m$  is even and  $n$  is odd in  $\int \sin^m x \cos^n x dx$ , we write  $n$  as  $2k + 1$  and use the identity  $\cos^2 x = 1 - \sin^2 x$  to obtain

$$\cos^n x = \cos^{2k+1} x = (\cos^2 x)^k \cos x = (1 - \sin^2 x)^k \cos x.$$

We then combine the single  $\cos x$  with  $dx$  and set  $\cos x dx$  equal to  $d(\sin x)$ .

**Case 3** If both  $m$  and  $n$  are even in  $\int \sin^m x \cos^n x dx$ , we substitute


$$\sin^2 x = \frac{1 - \cos 2x}{2}, \quad \cos^2 x = \frac{1 + \cos 2x}{2} \quad (2)$$

to reduce the integrand to one in lower powers of  $\cos 2x$ .






# 8.5


## Trigonometric Substitutions



**FIGURE 8.2** Reference triangles for the three basic substitutions identifying the sides labeled  $x$  and  $a$  for each substitution.



**FIGURE 8.3** The arctangent, arcsine, and arcsecant of  $x/a$ , graphed as functions of  $x/a$ .




**FIGURE 8.4** Reference triangle for  $x = 2 \tan \theta$  (Example 1):

$$\tan \theta = \frac{x}{2}$$

and

$$\sec \theta = \frac{\sqrt{4 + x^2}}{2}.$$




**FIGURE 8.5** Reference triangle for  $x = 3 \sin \theta$  (Example 2):


$$\sin \theta = \frac{x}{3}$$

and


$$\cos \theta = \frac{\sqrt{9 - x^2}}{3}.$$



**FIGURE 8.6** If  $x = (2/5)\sec \theta$ ,  $0 < \theta < \pi/2$ , then  $\theta = \sec^{-1}(5x/2)$ , and we can read the values of the other trigonometric functions of  $\theta$  from this right triangle (Example 3).




**FIGURE 8.7** The region (a) and solid (b) in Example 4.



**FIGURE 8.8** Reference triangle for  $x = 2 \tan \theta$  (Example 4).





**FIGURE 8.9** The ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  in

Example 5.

# 8.6

## Integral Tables and Computer Algebra Systems

**EXAMPLE 1** Find

$$\int x(2x + 5)^{-1} dx.$$

**Solution** We use Formula 8 (not 7, which requires  $n \neq -1$ ):

$$\int x(ax + b)^{-1} dx = \frac{x}{a} - \frac{b}{a^2} \ln |ax + b| + C.$$

With  $a = 2$  and  $b = 5$ , we have

$$\int x(2x + 5)^{-1} dx = \frac{x}{2} - \frac{5}{4} \ln |2x + 5| + C.$$

**EXAMPLE 2** Find

$$\int \frac{dx}{x\sqrt{2x+4}}.$$

**Solution** We use Formula 13(b):

$$\int \frac{dx}{x\sqrt{ax+b}} = \frac{1}{\sqrt{b}} \ln \left| \frac{\sqrt{ax+b} - \sqrt{b}}{\sqrt{ax+b} + \sqrt{b}} \right| + C, \quad \text{if } b > 0.$$

With  $a = 2$  and  $b = 4$ , we have

$$\begin{aligned} \int \frac{dx}{x\sqrt{2x+4}} &= \frac{1}{\sqrt{4}} \ln \left| \frac{\sqrt{2x+4} - \sqrt{4}}{\sqrt{2x+4} + \sqrt{4}} \right| + C \\ &= \frac{1}{2} \ln \left| \frac{\sqrt{2x+4} - 2}{\sqrt{2x+4} + 2} \right| + C. \end{aligned}$$

**EXAMPLE 3** Find

$$\int \frac{dx}{x\sqrt{2x-4}}.$$

**Solution** We use Formula 13(a):

$$\int \frac{dx}{x\sqrt{ax-b}} = \frac{2}{\sqrt{b}} \tan^{-1} \sqrt{\frac{ax-b}{b}} + C.$$

With  $a = 2$  and  $b = 4$ , we have

$$\int \frac{dx}{x\sqrt{2x-4}} = \frac{2}{\sqrt{4}} \tan^{-1} \sqrt{\frac{2x-4}{4}} + C = \tan^{-1} \sqrt{\frac{x-2}{2}} + C.$$

**EXAMPLE 4** Find

$$\int \frac{dx}{x^2 \sqrt{2x - 4}}.$$

**Solution** We begin with Formula 15:

$$\int \frac{dx}{x^2 \sqrt{ax + b}} = -\frac{\sqrt{ax + b}}{bx} - \frac{a}{2b} \int \frac{dx}{x \sqrt{ax + b}} + C.$$

With  $a = 2$  and  $b = -4$ , we have

$$\int \frac{dx}{x^2 \sqrt{2x - 4}} = -\frac{\sqrt{2x - 4}}{-4x} + \frac{2}{2 \cdot 4} \int \frac{dx}{x \sqrt{2x - 4}} + C.$$

We then use Formula 13(a) to evaluate the integral on the right (Example 3) to obtain

$$\int \frac{dx}{x^2 \sqrt{2x - 4}} = \frac{\sqrt{2x - 4}}{4x} + \frac{1}{4} \tan^{-1} \sqrt{\frac{x - 2}{2}} + C.$$

**EXAMPLE 5** Find

$$\int x \sin^{-1} x \, dx.$$

**Solution** We use Formula 99:

$$\int x^n \sin^{-1} ax \, dx = \frac{x^{n+1}}{n+1} \sin^{-1} ax - \frac{a}{n+1} \int \frac{x^{n+1} dx}{\sqrt{1-a^2x^2}}, \quad n \neq -1.$$

With  $n = 1$  and  $a = 1$ , we have

$$\int x \sin^{-1} x \, dx = \frac{x^2}{2} \sin^{-1} x - \frac{1}{2} \int \frac{x^2 dx}{\sqrt{1-x^2}}.$$

The integral on the right is found in the table as Formula 33:

$$\int \frac{x^2}{\sqrt{a^2-x^2}} dx = \frac{a^2}{2} \sin^{-1} \left( \frac{x}{a} \right) - \frac{1}{2} x \sqrt{a^2-x^2} + C.$$

With  $a = 1$ ,

$$\int \frac{x^2 dx}{\sqrt{1-x^2}} = \frac{1}{2} \sin^{-1} x - \frac{1}{2} x \sqrt{1-x^2} + C.$$

The combined result is

$$\begin{aligned} \int x \sin^{-1} x \, dx &= \frac{x^2}{2} \sin^{-1} x - \frac{1}{2} \left( \frac{1}{2} \sin^{-1} x - \frac{1}{2} x \sqrt{1-x^2} + C \right) \\ &= \left( \frac{x^2}{2} - \frac{1}{4} \right) \sin^{-1} x + \frac{1}{4} x \sqrt{1-x^2} + C'. \end{aligned}$$

**EXAMPLE 6** Using a Reduction Formula

Find

$$\int \tan^5 x \, dx.$$

**Solution** We apply Equation (1) with  $n = 5$  to get

$$\int \tan^5 x \, dx = \frac{1}{4} \tan^4 x - \int \tan^3 x \, dx.$$

We then apply Equation (1) again, with  $n = 3$ , to evaluate the remaining integral:

$$\int \tan^3 x \, dx = \frac{1}{2} \tan^2 x - \int \tan x \, dx = \frac{1}{2} \tan^2 x + \ln |\cos x| + C.$$

The combined result is

$$\int \tan^5 x \, dx = \frac{1}{4} \tan^4 x - \frac{1}{2} \tan^2 x - \ln |\cos x| + C'.$$



### EXAMPLE 7 Deriving a Reduction Formula

Show that for any positive integer  $n$ ,

$$\int (\ln x)^n dx = x(\ln x)^n - n \int (\ln x)^{n-1} dx.$$

**Solution** We use the integration by parts formula

$$\int u dv = uv - \int v du$$

with

$$u = (\ln x)^n, \quad du = n(\ln x)^{n-1} \frac{dx}{x}, \quad dv = dx, \quad v = x,$$

to obtain

$$\int (\ln x)^n dx = x(\ln x)^n - n \int (\ln x)^{n-1} dx.$$

**EXAMPLE 8** Find

$$\int \sin^2 x \cos^3 x \, dx.$$

**Solution 1** We apply Equation (3) with  $n = 2$  and  $m = 3$  to get

$$\begin{aligned} \int \sin^2 x \cos^3 x \, dx &= -\frac{\sin x \cos^4 x}{2 + 3} + \frac{1}{2 + 3} \int \sin^0 x \cos^3 x \, dx \\ &= -\frac{\sin x \cos^4 x}{5} + \frac{1}{5} \int \cos^3 x \, dx. \end{aligned}$$

We can evaluate the remaining integral with Formula 61 (another reduction formula):

$$\int \cos^n ax \, dx = \frac{\cos^{n-1} ax \sin ax}{na} + \frac{n-1}{n} \int \cos^{n-2} ax \, dx.$$

**Continued on next slide**

With  $n = 3$  and  $a = 1$ , we have

$$\begin{aligned}\int \cos^3 x \, dx &= \frac{\cos^2 x \sin x}{3} + \frac{2}{3} \int \cos x \, dx \\ &= \frac{\cos^2 x \sin x}{3} + \frac{2}{3} \sin x + C.\end{aligned}$$

The combined result is

$$\begin{aligned}\int \sin^2 x \cos^3 x \, dx &= -\frac{\sin x \cos^4 x}{5} + \frac{1}{5} \left( \frac{\cos^2 x \sin x}{3} + \frac{2}{3} \sin x + C \right) \\ &= -\frac{\sin x \cos^4 x}{5} + \frac{\cos^2 x \sin x}{15} + \frac{2}{15} \sin x + C' .\end{aligned}$$

**Solution 2** Equation (3) corresponds to Formula 68 in the table, but there is another formula we might use, namely Formula 69. With  $a = 1$ , Formula 69 gives

$$\int \sin^n x \cos^m x \, dx = \frac{\sin^{n+1} x \cos^{m-1} x}{m+n} + \frac{m-1}{m+n} \int \sin^n x \cos^{m-2} x \, dx.$$

In our case,  $n = 2$  and  $m = 3$ , so that


$$\begin{aligned}\int \sin^2 x \cos^3 x \, dx &= \frac{\sin^3 x \cos^2 x}{5} + \frac{2}{5} \int \sin^2 x \cos x \, dx \\ &= \frac{\sin^3 x \cos^2 x}{5} + \frac{2}{5} \left( \frac{\sin^3 x}{3} \right) + C \\ &= \frac{\sin^3 x \cos^2 x}{5} + \frac{2}{15} \sin^3 x + C.\end{aligned}$$

As you can see, it is faster to use Formula 69, but we often cannot tell beforehand how things will work out. Do not spend a lot of time looking for the “best” formula. Just find one that will work and forge ahead.

Notice also that Formulas 68 (Solution 1) and 69 (Solution 2) lead to different-looking answers. That is often the case with trigonometric integrals and is no cause for concern. The results are equivalent, and we may use whichever one we please.

# 8.7

## Numerical Integration



**FIGURE 8.10** The Trapezoidal Rule approximates short stretches of the curve  $y = f(x)$  with line segments. To approximate the integral of  $f$  from  $a$  to  $b$ , we add the areas of the trapezoids made by joining the ends of the segments to the  $x$ -axis.


## The Trapezoidal Rule

To approximate  $\int_a^b f(x) dx$ , use

$$T = \frac{\Delta x}{2} \left( y_0 + 2y_1 + 2y_2 + \cdots + 2y_{n-1} + y_n \right).$$

The  $y$ 's are the values of  $f$  at the partition points

$x_0 = a, x_1 = a + \Delta x, x_2 = a + 2\Delta x, \dots, x_{n-1} = a + (n - 1)\Delta x, x_n = b,$   
where  $\Delta x = (b - a)/n$ .



**TABLE 8.3**

$x$	$y = x^2$
1	1
$\frac{5}{4}$	$\frac{25}{16}$
$\frac{6}{4}$	$\frac{36}{16}$
$\frac{7}{4}$	$\frac{49}{16}$
2	4


**FIGURE 8.11** The trapezoidal approximation of the area under the graph of  $y = x^2$  from  $x = 1$  to  $x = 2$  is a slight overestimate (Example 1).

### The Error Estimate for the Trapezoidal Rule


If  $f''$  is continuous and  $M$  is any upper bound for the values of  $|f''|$  on  $[a, b]$ , then the error  $E_T$  in the trapezoidal approximation of the integral of  $f$  from  $a$  to  $b$  for  $n$  steps satisfies the inequality

$$|E_T| \leq \frac{M(b-a)^3}{12n^2}.$$







**FIGURE 8.12** Graph of the integrand in Example 3.



**FIGURE 8.13** The continuous function  $y = 2/x^3$  has its maximum value on  $[1, 2]$  at  $x = 1$ .



**FIGURE 8.14** Simpson's Rule approximates short stretches of the curve with parabolas.



**FIGURE 8.15** By integrating from  $-h$  to  $h$ , we find the shaded area to be

$$\frac{h}{3}(y_0 + 4y_1 + y_2).$$

## Simpson's Rule

To approximate  $\int_a^b f(x) dx$ , use

$$S = \frac{\Delta x}{3} (y_0 + 4y_1 + 2y_2 + 4y_3 + \cdots + 2y_{n-2} + 4y_{n-1} + y_n).$$

The  $y$ 's are the values of  $f$  at the partition points

$$x_0 = a, x_1 = a + \Delta x, x_2 = a + 2\Delta x, \dots, x_{n-1} = a + (n - 1)\Delta x, x_n = b.$$

The number  $n$  is even, and  $\Delta x = (b - a)/n$ .

**TABLE 8.4**

$x$	$y = 5x^4$
0	0
$\frac{1}{2}$	$\frac{5}{16}$
1	5
$\frac{3}{2}$	$\frac{405}{16}$
2	80

**EXAMPLE 5** Applying Simpson's Rule

Use Simpson's Rule with  $n = 4$  to approximate  $\int_0^2 5x^4 dx$ .

**Solution** Partition  $[0, 2]$  into four subintervals and evaluate  $y = 5x^4$  at the partition points (Table 8.4). Then apply Simpson's Rule with  $n = 4$  and  $\Delta x = 1/2$ :

$$\begin{aligned} S &= \frac{\Delta x}{3} \left( y_0 + 4y_1 + 2y_2 + 4y_3 + y_4 \right) \\ &= \frac{1}{6} \left( 0 + 4\left(\frac{5}{16}\right) + 2(5) + 4\left(\frac{405}{16}\right) + 80 \right) \\ &= 32\frac{1}{12}. \end{aligned}$$

This estimate differs from the exact value (32) by only  $1/12$ , a percentage error of less than three-tenths of one percent, and this was with just four subintervals.

### **The Error Estimate for Simpson's Rule**


If  $f^{(4)}$  is continuous and  $M$  is any upper bound for the values of  $|f^{(4)}|$  on  $[a, b]$ , then the error  $E_S$  in the Simpson's Rule approximation of the integral of  $f$  from  $a$  to  $b$  for  $n$  steps satisfies the inequality

$$|E_S| \leq \frac{M(b - a)^5}{180n^4}.$$

**TABLE 8.5** Trapezoidal Rule approximations ( $T_n$ ) and Simpson's Rule approximations ( $S_n$ ) of  $\ln 2 = \int_1^2 (1/x) dx$

$n$	$T_n$	Error  less than ...	$S_n$	Error  less than ...
10	0.6937714032	0.0006242227	0.6931502307	0.0000030502
20	0.6933033818	0.0001562013	0.6931473747	0.0000001942
30	0.6932166154	0.0000694349	0.6931472190	0.0000000385
40	0.6931862400	0.0000390595	0.6931471927	0.0000000122
50	0.6931721793	0.0000249988	0.6931471856	0.0000000050
100	0.6931534305	0.0000062500	0.6931471809	0.0000000004








**FIGURE 8.16** The dimensions of the swamp in Example 9.

# 8.8


## Improper Integrals



**FIGURE 8.17** Are the areas under these infinite curves finite?



(a)



(b)

**FIGURE 8.18** (a) The area in the first quadrant under the curve  $y = e^{-x/2}$  is (b) an improper integral of the first type.

## DEFINITION Type I Improper Integrals

Integrals with infinite limits of integration are **improper integrals of Type I**.

1. If  $f(x)$  is continuous on  $[a, \infty)$ , then

$$\int_a^{\infty} f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx.$$

2. If  $f(x)$  is continuous on  $(-\infty, b]$ , then


$$\int_{-\infty}^b f(x) dx = \lim_{a \rightarrow -\infty} \int_a^b f(x) dx.$$

3. If  $f(x)$  is continuous on  $(-\infty, \infty)$ , then


$$\int_{-\infty}^{\infty} f(x) dx = \int_{-\infty}^c f(x) dx + \int_c^{\infty} f(x) dx,$$

where  $c$  is any real number.

In each case, if the limit is finite we say that the improper integral **converges** and that the limit is the **value** of the improper integral. If the limit fails to exist, the improper integral **diverges**.



**FIGURE 8.19** The area under this curve is an improper integral (Example 1).



NOT TO SCALE

**FIGURE 8.20** The area under this curve is finite (Example 2).

### EXAMPLE 3 Determining Convergence

For what values of  $p$  does the integral  $\int_1^\infty dx/x^p$  converge? When the integral does converge, what is its value?

**Solution** If  $p \neq 1$ ,

$$\int_1^b \frac{dx}{x^p} = \left. \frac{x^{-p+1}}{-p+1} \right]_1^b = \frac{1}{1-p} (b^{-p+1} - 1) = \frac{1}{1-p} \left( \frac{1}{b^{p-1}} - 1 \right).$$

Thus,

$$\begin{aligned} \int_1^\infty \frac{dx}{x^p} &= \lim_{b \rightarrow \infty} \int_1^b \frac{dx}{x^p} \\ &= \lim_{b \rightarrow \infty} \left[ \frac{1}{1-p} \left( \frac{1}{b^{p-1}} - 1 \right) \right] = \begin{cases} \frac{1}{p-1}, & p > 1 \\ \infty, & p < 1 \end{cases} \end{aligned}$$

because


$$\lim_{b \rightarrow \infty} \frac{1}{b^{p-1}} = \begin{cases} 0, & p > 1 \\ \infty, & p < 1. \end{cases}$$

Therefore, the integral converges to the value  $1/(p-1)$  if  $p > 1$  and it diverges if  $p < 1$ .

If  $p = 1$ , the integral also diverges:

$$\begin{aligned} \int_1^\infty \frac{dx}{x^p} &= \int_1^\infty \frac{dx}{x} \\ &= \lim_{b \rightarrow \infty} \int_1^b \frac{dx}{x} \\ &= \lim_{b \rightarrow \infty} \ln x \Big|_1^b \\ &= \lim_{b \rightarrow \infty} (\ln b - \ln 1) = \infty. \end{aligned}$$





**FIGURE 8.21** The area under this curve is

$$\lim_{a \rightarrow 0^+} \int_a^1 \left( \frac{1}{\sqrt{x}} \right) dx = 2,$$

an improper integral of the second kind.

## DEFINITION Type II Improper Integrals

Integrals of functions that become infinite at a point within the interval of integration are **improper integrals of Type II**.

1. If  $f(x)$  is continuous on  $(a, b]$  and is discontinuous at  $a$  then

$$\int_a^b f(x) dx = \lim_{c \rightarrow a^+} \int_c^b f(x) dx.$$


2. If  $f(x)$  is continuous on  $[a, b)$  and is discontinuous at  $b$ , then

$$\int_a^b f(x) dx = \lim_{c \rightarrow b^-} \int_a^c f(x) dx.$$

3. If  $f(x)$  is discontinuous at  $c$ , where  $a < c < b$ , and continuous on  $[a, c) \cup (c, b]$ , then

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$


In each case, if the limit is finite we say the improper integral **converges** and that the limit is the **value** of the improper integral. If the limit does not exist, the integral **diverges**.



**FIGURE 8.22** The limit does not exist:

$$\int_0^1 \left( \frac{1}{1-x} \right) dx = \lim_{b \rightarrow 1^-} \int_0^b \frac{1}{1-x} dx = \infty$$


The area beneath the curve and above the  $x$ -axis for  $[0, 1)$  is not a real number (Example 4).




**FIGURE 8.23** Example 5 shows the convergence of

$$\int_0^3 \frac{1}{(x-1)^{2/3}} dx = 3 + 3\sqrt[3]{2},$$

so the area under the curve exists (so it is a real number).



**FIGURE 8.24** The calculation in Example 7 shows that this infinite horn has a finite volume.



**FIGURE 8.25** The graph of  $e^{-x^2}$  lies below the graph of  $e^{-x}$  for  $x > 1$  (Example 9).

### **THEOREM 1**    **Direct Comparison Test**

Let  $f$  and  $g$  be continuous on  $[a, \infty)$  with  $0 \leq f(x) \leq g(x)$  for all  $x \geq a$ . Then

1.  $\int_a^{\infty} f(x) dx$     converges if     $\int_a^{\infty} g(x) dx$     converges

2.  $\int_a^{\infty} g(x) dx$     diverges if     $\int_a^{\infty} f(x) dx$     diverges.

## THEOREM 2    Limit Comparison Test

If the positive functions  $f$  and  $g$  are continuous on  $[a, \infty)$  and if


$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = L, \quad 0 < L < \infty,$$

then

$$\int_a^{\infty} f(x) \, dx \quad \text{and} \quad \int_a^{\infty} g(x) \, dx$$

both converge or both diverge.






**FIGURE 8.26** The functions in Example 11.

## Types of Improper Integrals Discussed in This Section

INFINITE LIMITS OF INTEGRATION: **TYPE I**


### 1. Upper limit

$$\int_1^{\infty} \frac{\ln x}{x^2} dx = \lim_{b \rightarrow \infty} \int_1^b \frac{\ln x}{x^2} dx$$




### 2. Lower limit

$$\int_{-\infty}^0 \frac{dx}{1+x^2} = \lim_{a \rightarrow -\infty} \int_a^0 \frac{dx}{1+x^2}$$



### 3. Both limits


$$\int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \lim_{b \rightarrow -\infty} \int_b^0 \frac{dx}{1+x^2} + \lim_{c \rightarrow \infty} \int_0^c \frac{dx}{1+x^2}$$



INTEGRAND BECOMES INFINITE: **TYPE II**


### 4. Upper endpoint

$$\int_0^1 \frac{dx}{(x-1)^{2/3}} = \lim_{b \rightarrow 1^-} \int_0^b \frac{dx}{(x-1)^{2/3}}$$




### 5. Lower endpoint


$$\int_1^3 \frac{dx}{(x-1)^{2/3}} = \lim_{d \rightarrow 1^+} \int_d^3 \frac{dx}{(x-1)^{2/3}}$$



### 6. Interior point

$$\int_0^3 \frac{dx}{(x-1)^{2/3}} = \int_0^1 \frac{dx}{(x-1)^{2/3}} + \int_1^3 \frac{dx}{(x-1)^{2/3}}$$





**FIGURE 8.27** Euler's gamma function  $\Gamma(x)$  is a continuous function of  $x$  whose value at each positive integer  $n + 1$  is  $n!$ . The defining integral formula for  $\Gamma$  is valid only for  $x > 0$ , but we can extend  $\Gamma$  to negative noninteger values of  $x$  with the formula  $\Gamma(x) = (\Gamma(x + 1))/x$ , which is the subject of Exercise 49.